

# STAT 3743: Probability and Statistics

G. Jay Kerns, Youngstown State University

Fall 2010

G. Jay Kerns, Youngstown State University

[Probability and Statistics](#)



Notes

---

---

---

---

---

---

---

---

---

---

## Probability

- Random experiment: outcome not known in advance
- Sample space: set of all possible outcomes ( $S$ )
- Probability related to Set Theory
  - subsets  $A, B, C$ , etc. are *events*
  - $\emptyset$  represents the empty set

### How to do it with R

```
> library(prob)
> S <- data.frame(lands = c("down",
+   "up", "side"))
> S <- tosscoin(3)
```

G. Jay Kerns, Youngstown State University

[Probability and Statistics](#)



Notes

---

---

---

---

---

---

---

---

---

---

## Set Theory review

Name	Denoted	Defined by elements	R syntax
Union	$A \cup B$	in $A$ or $B$ or both	<code>union(A, B)</code>
Intersection	$A \cap B$	in both $A$ and $B$	<code>intersect(A, B)</code>
Difference	$A \setminus B$	in $A$ but not in $B$	<code>setdiff(A, B)</code>
Complement	$A^c$	in $S$ but not in $A$	<code>setdiff(S, A)</code>

Table: Set operations

Notes

---

---

---

---

---

---

---

---

## Algebra of sets

- $A \cup \emptyset = A$ ,  $A \cap \emptyset = \emptyset$ ,  $A \cup S = S$ ,  $A \cap S = A$ , ...
- Commutative property:

$$A \cup B = B \cup A, \quad A \cap B = B \cap A$$

- Associative property:

$$(A \cup B) \cup C = A \cup (B \cup C), \quad (A \cap B) \cap C = A \cap (B \cap C)$$

- Distributive property:

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C), \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

Notes

---

---

---

---

---

---

---

---

### Example

Write “neither  $A$  nor  $B$  occurs”

### Example

“ $A$  occurs, but not  $B$ ”

### Example

“ $A$  or  $B$  occurs, but not both”



### Notes

---

---

---

---

---

---

---

---

---

---

### Definition

The sets  $A$  and  $B$  are *mutually exclusive* or *disjoint* if  $A \cap B = \emptyset$ . We say  $A_1, A_2, \dots, A_k$  are m.e. if  $A_i \cap A_j = \emptyset$  when  $i \neq j$ .

- Have all kinds of events, want to know chance of an event  $A$
- The probability of  $A$  is the proportion of times that  $A$  occurs in repeated trials of a random experiment as the number of trials increases without bound.



### Notes

---

---

---

---

---

---

---

---

---

---

# Axioms for Probability

## Axiom 1.

$\mathbb{P}(A) \geq 0$  for any event  $A \subset S$ .

## Axiom 2.

$\mathbb{P}(S) = 1$ .

## Axiom 3.

If the events  $A_1, A_2, A_3, \dots$  are disjoint then

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i).$$

Navigation icons: back, forward, search, etc.

Notes

---

---

---

---

---

---

---

---

---

---

# Properties of probability

## Property 1.

$\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$

## Property 2.

$\mathbb{P}(\emptyset) = 0$

## Property 3.

If  $A \subset B$ , then  $\mathbb{P}(A) \leq \mathbb{P}(B)$

Navigation icons: back, forward, search, etc.

Notes

---

---

---

---

---

---

---

---

---

---

## Properties of probability

### Property 4.

$$0 \leq \mathbb{P}(A) \leq 1$$

### Property 5. (General Addition Rule)

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$$



Notes

---

---

---

---

---

---

---

---

---

---

## Properties of probability

### What about 3 events?

### Corollary. (Boole's Inequality)

$$\mathbb{P}(A \cup B) \leq \mathbb{P}(A) + \mathbb{P}(B)$$



Notes

---

---

---

---

---

---

---

---

---

---

## How do we assign probabilities?

Finite sample space

$$S = \{e_1, e_2, \dots, e_N\}$$

Need

- ①  $p_i \geq 0$
- ②  $\mathbb{P}(S) = \sum_{i=1}^N p_i = 1$

**Equally likely outcomes**

means

$$p_1 = p_2 = \dots = p_N = p \implies p = 1/N$$

Notes

---

---

---

---

---

---

---

---

---

---

## How do we assign probabilities?

Given  $A \subset S$ , write

$$A = \{a_{i_1}, a_{i_2}, \dots, a_{i_k}\}$$

Then

$$\begin{aligned}\mathbb{P}(A) &= \mathbb{P}(a_{i_1}) + \mathbb{P}(a_{i_2}) + \dots + \mathbb{P}(a_{i_k}), \\ &= \frac{1}{N} + \frac{1}{N} + \dots + \frac{1}{N}, \\ &= \frac{k}{N} = \frac{\#(A)}{\#(S)}.\end{aligned}$$

Notes

---

---

---

---

---

---

---

---

---

---

## Examples

### Example 1.

Toss a coin

### Example 2.

Toss 2 coins

$$\mathbb{P}(\text{at least 1 head}) =$$
$$\mathbb{P}(\text{no heads}) =$$

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

G. Jay Kerns, Youngstown State University

## Probability and Statistics

## Notes

[illegible]

## Examples

### Example 3.

Three child family

$$\mathbb{P}(\text{exactly 2 boys}) =$$
$$\mathbb{P}(\text{at most 2 boys}) =$$

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ 🔍 ↺

G. Jay Kerns, Youngstown State University

## Probability and Statistics

## Notes

[illegible]

## Examples

### Example 5.

Deck of cards. Select 1 card at random.

$$A \longrightarrow \{\text{Ace}\} \quad \mathbb{P}(A) =$$

$$B \longrightarrow \{\text{Clubs}\} \quad \mathbb{P}(B) =$$

$$\mathbb{P}(A \cap B) =$$

$$\mathbb{P}(A \cup B) =$$



Notes

---

---

---

---

---

---

---

---

---

---

## Examples

### Example 6.

Poker hand  $\longrightarrow$  STUD poker

$$S = \{ \quad \quad \quad \}$$

$$\mathbb{P}(\text{Royal Flush}) =$$



Notes

---

---

---

---

---

---

---

---

---

---

# How to count

## Multiplication Principle.

An experiment has two steps. First step can be done in  $n_1$  ways, Second step can be done in  $n_2$  ways. The whole experiment may be done in

$$n_1 n_2 \text{ ways}$$

If it has  $k$  steps which can be done in  $n_1, n_2, \dots, n_k$  ways, then the whole experiment may be done in

$$n_1 n_2 \cdots n_k \text{ ways}$$

Notes

---

---

---

---

---

---

---

---

---

---

# Examples

## Examples.

① Want to eat a pizza

② Toss 6 coins

③ Roll 112 dice

What about  $\mathbb{P}(70 \text{ sixes})$ ?

Notes

---

---

---

---

---

---

---

---

---

---

# How to count

## Theorem.

The number of ways to select an *ordered* sample of  $k$  subjects from a population that has  $n$  distinguishable members is

- $n^k$  if sampling is done with replacement,
- $n(n-1)(n-2)\cdots(n-k+1)$  if sampling is done without replacement.

Here, ORDER is IMPORTANT



Notes

---

---

---

---

---

---

---

---

---

---

# Examples

## Examples.

- 1 Flip a coin 7 times
- 2 20 students, select president, vice-president, treasurer
- 3 Rent 5 movies. Want to watch 3 movies on the first night.



Notes

---

---

---

---

---

---

---

---

---

---

## How to count

### Theorem.

The number of ways to select an *unordered* sample of  $k$  subjects from a population that has  $n$  distinguishable members is

- $(n - 1 + k)! / [(n - 1)!k!]$  if sampling is done with replacement,
- $n! / [k!(n - k)!]$  if sampling is done without replacement.

$n! / [k!(n - k)!]$  is a *binomial coefficient* “ $n$  choose  $k$ ”

$$\binom{n}{k} = \frac{n!}{k!(n - k)!}$$

Notes

---

---

---

---

---

---

---

---

---

---

## More about binomial coefficients

Notes

---

---

---

---

---

---

---

---

---

---

# Birthday problem

- $n$  people in a class
- 365 days/year, equally likely
- $\mathbb{P}(\text{at least two have same birthday})$

$$1 - \frac{\#(A)}{\#(S)} =$$

Notes

---

---

---

---

---

---

---

---

---

---




Figure: The birthday problem

Notes

---

---

---

---

---

---

---

---

---

---

## Poker hands

52 cards  $\longrightarrow$  5 card hand

$$S = \{\text{all possible 5 card hands}\}$$

(should shuffle times???)

$$A = \text{Royal Flush} = \{A, K, Q, J, \text{all same suit}\}$$

$$B = \{\text{Four of a kind}\}$$

Notes

---

---

---

---

---

---

---

---

---

---

## Conditional probability

52 cards  $\longrightarrow$  draw 2 cards (without replacement)

$$A = \{\text{1st card drawn is Ace}\}$$

$$B = \{\text{2nd card drawn is Ace}\}$$

Then

$$\mathbb{P}(A) =$$

$$\mathbb{P}(B) = \left\{ \right.$$

Notes

---

---

---

---

---

---

---

---

---

---

## Conditional probability

### Definition.

The conditional probability of  $B$  given that the event  $A$  occurred is

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)}, \quad \text{if } \mathbb{P}(A) > 0.$$

Notes

---

---

---

---

---

---

---

---

---

---

## Conditional probability

### Example

Toss a coin twice.

$$A = \{\text{a head occurs}\}$$

$$B = \{\text{a head and tail occurs}\}$$

- $\mathbb{P}(A|B) =$
- $\mathbb{P}(B|A) =$

Notes

---

---

---

---

---

---

---

---

---

---

# Conditional probability

Example

Toss a die twice.

$$A = \{\text{outcomes match}\}$$

$$B = \{\text{sum of outcomes} \geq 8\}$$

- $\mathbb{P}(A) =$
- $\mathbb{P}(B) =$
- $\mathbb{P}(A \cap B) =$
- $\mathbb{P}(A|B) =$
- $\mathbb{P}(B|A) =$

Notes

---

---

---

---

---

---

---

---

---

---

# Properties

**Note.**

For any fixed event  $A$  with  $\mathbb{P}(A) > 0$ ,

- 1  $\mathbb{P}(B|A) \geq 0$ , for all events  $B \subset S$ ,
- 2  $\mathbb{P}(S|A) = 1$ , and
- 3 If  $B_1, B_2, B_3, \dots$  are disjoint events, then

$$\mathbb{P}\left(\bigcup_{k=1}^{\infty} B_k \mid A\right) = \sum_{k=1}^{\infty} \mathbb{P}(B_k|A).$$

Notes

---

---

---

---

---

---

---

---

---

---

## More properties

### Note.

For any events  $A$ ,  $B$ , and  $C$  with  $\mathbb{P}(A) > 0$ ,

- ①  $\mathbb{P}(B^c|A) = 1 - \mathbb{P}(B|A)$ .
- ② If  $B \subset C$  then  $\mathbb{P}(B|A) \leq \mathbb{P}(C|A)$ .
- ③  $\mathbb{P}[(B \cup C)|A] = \mathbb{P}(B|A) + \mathbb{P}(C|A) - \mathbb{P}[(B \cap C)|A]$ .
- ④ For any two events  $A$  and  $B$ ,

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \mathbb{P}(B|A).$$

For 3 events:



Notes

---

---

---

---

---

---

---

---

---

---

## Conditional probability

### Example.

Recall the aces problem

$$A = \{\text{1st card drawn is Ace}\}$$

$$B = \{\text{2nd card drawn is Ace}\}$$

- $\mathbb{P}(\text{both Aces}) =$



Notes

---

---

---

---

---

---

---

---

---

---

# Conditional probability

## Example.

Urn with 10 balls, 7 red and 3 green. Select 3 balls successively from the urn.

$$A = \{\text{1st ball red}\}$$

$$B = \{\text{2nd ball red}\}$$

$$C = \{\text{3rd ball red}\}$$

- $IP(\text{all red}) =$

Notes

---

---

---

---

---

---

---

---

---

---

# Good example

Two urns. First: 5 red, 3 green. Second: 2 red, 6 green  
1 ball transferred. Select 1 ball.

$$IP(\text{red}) =$$

Notes

---

---

---

---

---

---

---

---

---

---

## What if you don't look?

---

$IP(\text{second card is Ace}) =$

Notes

---

---

---

---

---

---

---

---

---

---

---

## Good example (continued)

---

$IP(\text{red}) =$

Notes

---

---

---

---

---

---

---

---

---

---

---

# What if you don't look?

$\mathbb{P}(\text{second card is Ace}) =$

Notes

---

---

---

---

---

---

---

---

---

---

# Independence

## Example.

Toss two coins

- $\mathbb{P}(\text{1st } H) =$
- $\mathbb{P}(\text{2nd } H) =$
- $\mathbb{P}(\text{both } H) =$

$\mathbb{P}(\text{2nd } H \mid \text{1st } H) =$

Notes

---

---

---

---

---

---

---

---

---

---

# Independence

**Definition.**

Events  $A$  and  $B$  are *independent* if

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \mathbb{P}(B),$$

otherwise they are *dependent*.

Intuition:

$$\mathbb{P}(A|B) = \mathbb{P}(A) \quad \text{when } A, B \text{ independent}$$

Notes

---

---

---

---

---

---

---

---

---

---

# Properties

**Proposition.**

If  $A$  and  $B$  are independent then

- $A$  and  $B^c$  are independent,
- $A^c$  and  $B$  are independent,
- $A^c$  and  $B^c$  are independent.

What about 3 or more events?

Notes

---

---

---

---

---

---

---

---

---

---

## Mutual independence

### Definition.

$A$ ,  $B$  and  $C$  are *mutually independent* if

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \mathbb{P}(B),$$

$$\mathbb{P}(A \cap C) = \mathbb{P}(A) \mathbb{P}(C),$$

$$\mathbb{P}(B \cap C) = \mathbb{P}(B) \mathbb{P}(C),$$

and

$$\mathbb{P}(A \cap B \cap C) = \mathbb{P}(A) \mathbb{P}(B) \mathbb{P}(C).$$



Notes

---

---

---

---

---

---

---

---

---

---

## Mutual independence

### Example.

Toss 100 coins.

$$\mathbb{P}(\text{at least 1 head}) =$$



Notes

---

---

---

---

---

---

---

---

---

---

# Mutual independence

## Remark.

Pairwise independence does NOT imply mutual.

## Examples.

- ① Toss coins, roll dice, *etc.*
- ② Draw two cards *without* replacement
- ③ Space shuttle. 4 computers,  $A, B, C, D$

$$\mathbb{P}(\text{fail}) = 0.10$$



Notes

---

---

---

---

---

---

---

---

---

---

# Space shuttle (cont.)

Scheme: computers in series.

If computers independent,

$$\mathbb{P}(\text{at least one computer works})$$



Notes

---

---

---

---

---

---

---

---

---

---

# Bayes' Rule

## Theorem.

Let  $B_1, B_2, \dots, B_n$  be mutually exclusive and exhaustive and let  $A$  be an event with  $\mathbb{P}(A) > 0$ . Then

$$\mathbb{P}(B_k|A) = \frac{\mathbb{P}(B_k) \mathbb{P}(A|B_k)}{\sum_{i=1}^n \mathbb{P}(B_i) \mathbb{P}(A|B_i)}, \quad k = 1, 2, \dots, n.$$



# Bayes' Rule (intuition)



Notes

---

---

---

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

---

---

---

## Bayes' Rule: what does it mean?

Given (or know) *a priori* probabilities  $\mathbb{P}(B_k)$ . Collect some data, which is  $A$ .

How to update  $\mathbb{P}(B_k)$  to  $\mathbb{P}(B_k|A)$ ?

G. Jay Kerns, Youngstown State University

[Probability and Statistics](#)



Notes

---

---

---

---

---

---

---

---

---

---

## Example: misfiling assistants

Moe, Larry, and Curly

	Moe	Larry	Curly
Workload	60%	30%	10%

	Moe	Larry	Curly
Prior	$\mathbb{P}(M) =$	$\mathbb{P}(L) =$	$\mathbb{P}(C) =$

G. Jay Kerns, Youngstown State University

[Probability and Statistics](#)



Notes

---

---

---

---

---

---

---

---

---

---

## Misfiling assistants (cont.)

	Moe	Larry	Curly
Misfile Rate	0.003	0.007	0.010

	Moe	Larry	Curly
Posterior	$\text{IP}(M A) \approx$	$\text{IP}(L A) \approx$	$\text{IP}(C A) \approx$

G. Jay Kerns, Youngstown State University

## Probability and Statistics

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

## Random variables

- Experiment  $E$
- Sample space  $S$
- Calculate number  $X$

**Definition.**

A *random variable*  $X$  is a function  $X : S \rightarrow \mathbb{R}$  that associates to each outcome  $\omega \in S$  exactly one number  $X(\omega) = x$ . The support of  $X$  is the set of  $X$ 's values:

$$S_X = \{x : X(\omega) = x, \omega \in S\}$$

G. Jay Kerns, Youngstown State University

## Probability and Statistics

## Notes

[illegible]

## Notes

[illegible]

# Random variables

**Example.**

Toss a coin three times

**Example.**

Toss a coin until tails

**Example.**

Toss a coin, measure time until lands



Notes

---

---

---

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

---

---

---